

## 2017 STD Surveillance Report Supplemental Data Sheet

	Chlamydia	Gonorrhea	P&S Syphilis	Congenital Syphilis
<b>Overall</b>	<ul style="list-style-type: none"> <li>Rate: 528.8 per 100,000</li> <li># of cases: 1,708,569 cases</li> <li>Change from 2016: 6.9% increase</li> </ul>	<ul style="list-style-type: none"> <li>Rate: 171.9 per 100,000</li> <li># of cases: 555,608</li> <li>Change from 2016: 18.6% increase</li> </ul>	<ul style="list-style-type: none"> <li>Rate: 9.5 per 100,000</li> <li># of cases: 30,644</li> <li>Change from 2016: 17.6% increase</li> </ul>	<ul style="list-style-type: none"> <li>Rate: 23.3 per 100,000 live births</li> <li># of cases: 918</li> <li>Change from 2016: 43.8% increase</li> </ul>
<b>Women</b>	<ul style="list-style-type: none"> <li>5.1% increase</li> <li>3<sup>rd</sup> year of increases in rates among 15-19 y.o. (6.5% increase)</li> </ul>	<ul style="list-style-type: none"> <li>17.8% increase</li> </ul>	<ul style="list-style-type: none"> <li>Increase of 21.1% among women</li> </ul>	
<b>Men</b>	<ul style="list-style-type: none"> <li>10.5% increase</li> <li>3<sup>rd</sup> year of increases in rates among 15-19 y.o. (11.1% increase)</li> </ul>	<ul style="list-style-type: none"> <li>19.3% increase</li> <li>Estimated gonorrhea rates among MSM in SSuN jurisdictions increased 155% from 2013-2017</li> </ul>	<ul style="list-style-type: none"> <li>9.0% increase</li> <li>Men account for 88% of all cases</li> <li>68% of P&amp;S syphilis cases (male cases with known sex of sex partner and female cases) were MSM</li> </ul>	

Screening Recommendations	
<b>Screening Recommendations</b>	<p><u>Women</u></p> <ul style="list-style-type: none"> <li>If you are a sexually active woman younger than 25, or have risk factors such as new or multiple sex partners, you should request annual chlamydia and gonorrhea tests.</li> <li>If you are a pregnant woman, you should request syphilis, HIV, chlamydia, and hepatitis B tests at your first prenatal visit. If you have new or multiple sex partners, you should also request gonorrhea testing early in pregnancy.</li> <li>If you are a pregnant woman at high risk for syphilis, you should also be tested for syphilis at the beginning of your 3<sup>rd</sup> trimester and again at delivery.</li> </ul> <p><u>MSM</u></p> <ul style="list-style-type: none"> <li>If you are a sexually active man who is gay, bisexual, or has sex with men, you should request tests for syphilis, chlamydia, gonorrhea, and HIV at least once a year. More frequent STD testing is recommended for men at high risk.</li> </ul>