

For HIV, STDs, and Hepatitis test after each partner or exposure, or every 1-3 months for frequent partners or exposures.

Wyoming Department of Health Communicable Disease Snapshot of Screening Recommendations

| <u>At Risk Populations</u> | <u>HIV</u> | <u>Hep B*</u> | <u>Hep C</u> | <u>Syphilis</u> | <u>Chlamydia</u> | <u>Gonorrhea</u> | <u>TB</u> |
|--|------------|---------------|--------------|-----------------|------------------|------------------|-----------|
| Sexually active individuals | X | | | | X | X | |
| **Genital chlamydia/gonorrhea: history of & never tested or history of genital sex since last tested | X | | | | X | X | |
| Pharyngeal chlamydia/gonorrhea: history of & never tested or history of oral sex since last tested | X | | | | X | X | |
| Rectal chlamydia/gonorrhea: history of & never tested or history of anal sex since last tested | X | | | | X | X | |
| New sex partner in the last 60 days | X | | | | X | X | |
| More than one sex partner in last 60 days | X | | | X | X | X | |
| Sexual contact with a STD positive individual (as indicated) | X | | | X | X | X | |
| *Men who have sex with men (MSM) | X | X | X | X | X | X | |
| History of prior sexually transmitted diseases (STD) | X | | | | X | X | |
| Pelvic inflammatory disease (PID) | X | | | | X | X | |
| *Pregnant women | X | X | | X | X | X | |
| Sexual assault | X | | X | X | X | X | |
| *Current or past history of injection or intranasal drug use | X | X | X | | | | X |
| Current or past history of homelessness | X | X | X | | | | X |
| Current or past resident of detention or correction facility | X | X | X | | X | X | X |
| *HIV infected | | X | X | X | X | X | X |
| 1945-1965 birth cohort (Baby Boomers) | X | | X | | | | |
| *Hepatitis B or C positive sexual contact (as indicated) | X | X | X | | X | X | |
| *Household contact to Hepatitis B positive individuals | X | X | | | | | |
| *Current or past long-term hemodialysis | X | X | X | | | | |
| *Recipient of blood transfusions, blood components, or organ transplant prior to July 1992 | X | X | X | | | | |
| *Recipient of clotting factor or blood concentrate prior to 1987 | X | X | X | | | | |
| *Consistently abnormal liver tests | | X | X | | | | |
| *Underlying immunocompromising conditions/immunosuppressive (including chemo) therapy | X | X | | | | | X |
| Born in Latin America, Caribbean, Africa, Eastern Europe, and Russia | X | | | | | | X |
| *Born in Asia, Africa or South America | X | X | | | | | X |
| *Parents born in Asia, Africa, or South America | X | X | | | | | |
| Active tuberculosis (TB) disease | X | | | | | | |
| Household contact of patients with active TB disease | | | | | | | X |
| Travelers with prolonged stays in countries where TB is endemic | | | | | | | X |
| As indicated by employment (e.g. healthcare or congregate settings) | | | | | | | X |
| *Symptoms consistent with disease (as indicated) | X | X | X | X | X | X | X |

*Prior to Hepatitis B testing check WyIR for vaccination status. Do not test fully vaccinated patients for Hepatitis B unless vaccine failure is suspected.

**For genital chlamydia/gonorrhea testing a vaginal self-swab is preferred for individuals with a vagina and urine is preferred for those without a vagina.